



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let figure 2 represent the force diagram, $ab=W$, $bc=R$, $ac=R_1$, $\angle abc=90^\circ-\theta$, $\angle bac=90^\circ-\varphi$, $\angle acb=\theta+\varphi$.

$$\therefore R=W\cos\varphi/\sin(\theta+\varphi), R_1=W\cos\theta/\sin(\theta+\varphi).$$

$$\therefore R=5.5078 \text{ pounds, } R_1=7.1881 \text{ pounds.}$$

DIOPHANTINE ANALYSIS.

61. Proposed by SYLVESTER ROBBINS, North Branch Depot, New Jersey.

Investigate that infinite series of prime, integral, rational scalene triangles where the sides of every term are consecutive numbers; then take the necessary factors from the proper KEY, and by an expeditious method, find in their order the areas of ten initial terms.

Solution by the PROPOSER.

I. The KEY to this series of rational triangles is $\sqrt{3}=1, \frac{2}{1}, \frac{5}{3}, \frac{7}{4}, \frac{19}{11}, \frac{26}{11}, \frac{71}{41}, \frac{97}{26}, \frac{265}{153}, \frac{362}{209}, \frac{971}{780}, \frac{1351}{2911}, \frac{5042}{2911}, \frac{13775}{7953}, \frac{18817}{10864}, \frac{51409}{29681}, \frac{70226}{40545}, \frac{191861}{110771}, \frac{262087}{151316}$, etc. Regard the mean side as the base, and drop perpendicular from the opposite angle. Let $x=\frac{1}{2}$ base. Notice that $x-2$ and $x+2$ are the segments of the base, and $\sqrt{[3(x^2-1^2)]}$ is the altitude of the triangle. Find such values for x as will render $\sqrt{[3(x^2-1^2)]}$ rational.

When $x=1, 2, 7, 26, 97, 362, 1351, 5042, 18817, 70226, 262087$, $\sqrt{[3(x^2-1^2)]}=0, 3, 12, 45, 168, 627, 2320, 8733, 32592, 121635, 453948$, etc.

These values of x are the half-bases of the several triangles. They are also the numerators of the even convergents in the expansion of $\sqrt{3}$. The values of $\sqrt{[3(x^2-1^2)]}$ are the altitudes of the same triangles, respectively, and they are also three times the denominators of the even convergents in the expansion of $\sqrt{3}$. Multiply one-half the base of a triangle by its perpendicular height, or, three times the product of the terms of the n th even convergent, must give the area of the n th triangle in the series.

Thus, $3 \times 2 \times 1 = 6$; $3 \times 7 \times 4 = 84$; $3 \times 26 \times 15 = 1170$; $3 \times 97 \times 56 = 16296$; $3 \times 362 \times 209 = 226974$; $3 \times 1351 \times 780 = 3161340$; $3 \times 5042 \times 2911 = 44031786$; $3 \times 18817 \times 10864 = 613283664$; $3 \times 70226 \times 40545 = 8541939510$; $3 \times 262087 \times 151316 = 118973869476$, etc.

II. Numerators of even convergents in expansion of $\sqrt{3}$: $1, 2, 7, 26, 97, 362, 1351, 5042, 18817, 70226$, etc. Then $\frac{1}{3}(7^2-1^2)=6$; $\frac{1}{3}(26^2-2^2)=84$; $\frac{1}{3}(97^2-7^2)=1170$; $\frac{1}{3}(362^2-26^2)=16296$; $\frac{1}{3}(1351^2-97^2)=226974$; etc.

III. Denominators of even convergents: $1, 4, 15, 56, 209, 780, 2911$, etc. $\frac{1}{3}(4^2-0)=6$; $\frac{1}{3}(15^2-1^2)=84$; $\frac{1}{3}(56^2-4^2)=1170$; $\frac{1}{3}(209^2-15^2)=16296$; $\frac{1}{3}(780^2-56^2)=226974$; etc.

IV. Let x =the half-sum of the three sides of the triangle. Then $\frac{1}{2}x-1$, $\frac{1}{2}x$ and $\frac{1}{2}x+1$ are the remainders.

$$(x)[(\frac{1}{2}x)-1](\frac{1}{2}x)[(\frac{1}{2}x)+1]=\text{square of triangle. } 3(x^2-3^2)=\text{square.}$$

$$\sqrt{\{(\frac{1}{2}x^2)[(x^2-3^2)/9]\}}/x=\frac{1}{3}\sqrt{[(x^2-3^2)/3]}, \text{ the radius of inscribed circle.}$$

$$\text{Put } x=y+6; \text{ then } 3[(y+6)^2-3^2]=\text{square}=(my+9)^2.$$

$$3y^2+36y+81=m^2y^2+18my+81; \quad 3y+36=m^2y+18m. \quad y=(18m-36)/(3-m^2); \text{ and } x=y+6=(18m-18-6m^2)/(3-m^2).$$